

**TETRASOL SOLUBLE POWDER- tetracycline hydrochloride powder, for solution**

**Dechra Veterinary Products**

-----  
**TETRASOL SOLUBLE POWDER**

(tetracycline hydrochloride soluble powder)

**Antibiotic**

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**FOR USE IN ANIMALS ONLY**

**KEEP OUT OF REACH OF CHILDREN**

**Each pound contains 324 g of tetracycline hydrochloride.**

**Approved by FDA under ANADA # 200-234**

**Net Weight: 5 lbs. (2.27 kg)**

**INDICATIONS:** For use in the control and treatment of the following conditions in swine, calves and poultry.

**SWINE AND CALVES**

**INDICATIONS:** Control and treatment of bacterial enteritis (scours) caused by *Escherichia coli*; bacterial pneumonia associated with *Actinobacillus pleuropneumoniae*, *Pasteurella* spp. and *Klebsiella* spp. sensitive to tetracycline hydrochloride.

**RECOMMENDED DOSAGE LEVEL:** Use soluble powder in the drinking water at a drug level of tetracycline hydrochloride per gallon to provide 10 mg/lb of body weight per day in divided doses.

**DIRECTIONS FOR USE:** Administer for 3-5 days.

**CAUTION:** Do not use for more than 5 consecutive days.

**CHICKENS**

**INDICATIONS:** Control of chronic respiratory disease (CRD) and air sac disease caused by *Mycoplasma gallisepticum* and *Escherichia coli*; infectious synovitis caused by *Mycoplasma synoviae* sensitive to tetracycline hydrochloride.

**RECOMMENDED DOSAGE LEVEL:** CRD and air sac disease: Use soluble powder in the drinking water at a drug level of 400-800 mg tetracycline hydrochloride per gallon. Infectious synovitis: Use soluble powder in the drinking water at a drug level of 200-400 mg tetracycline hydrochloride per gallon.

**TURKEYS**

**INDICATIONS:** Control of infectious synovitis caused by *Mycoplasma synoviae*; bluecomb (transmissible enteritis, coronaviral enteritis) complicated by organisms sensitive to tetracycline hydrochloride.

**RECOMMENDED DOSAGE LEVEL:** Infectious synovitis: Use soluble powder in the

drinking water at a drug level of 400 mg tetracycline hydrochloride per gallon. Bluecomb: Use soluble powder in the drinking water at a drug level of tetracycline hydrochloride per gallon to provide 25 mg/lb of body weight per day in divided doses.

**DIRECTIONS FOR USE:** Administer for 7-14 days. Medicate at first clinical signs of disease or when experience indicates the disease may be a problem.

**CAUTION:** Not for use in turkeys or chickens producing eggs for human consumption. Do not use for more than 14 consecutive days.

**WARNING:**

**SWINE** – Do not slaughter animals for food purposes within 4 days if treatment.

**CALVES** – Do not slaughter animals for food purposes within 5 days if treatment. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**CHICKENS and TURKEYS** - Do not slaughter birds for food within 4 days of treatment.

**GENERAL CAUTION:**

Prepare fresh solutions at least once a day. Solution are not stable for more than 24 hours. Use as sole source of tetracycline. Diagnosis should be reconsidered if improvement is not noticed within 3 days. The concentration of drug required in medicated water must be adequate to compensate for variations in the age and class of animals, feed consumption, and environmental temperature and humidity, each of which affects water consumption.

**MIXING DIRECTIONS**

**FOR SWINE, CALVES AND TURKEYS:** 2.52 oz. (two scoops) dissolved in 500 mL (approximately 17 fl. oz.) of warm water will provide a stock solution of 100 mg of tetracycline hydrochloride activity per mL.

**FOR TURKEYS ONLY:** This stock solution when metered at approximately 1 oz. per gallon, will provide drinking water containing 2,957 mg of tetracycline hydrochloride activity per gallon.

The contents of two scoops are sufficient to deliver the recommended daily dosage levels as follows:

At 25 mg/lb of body weight .... 2,040 total lbs of turkeys to be medicated.

At 10 mg/lb of body weight .... 5,100 total lbs of swine or calves to be medicated.

Individual calf treatment ..... 5 mL (1 measuring teaspoonful) twice daily for each 100 lbs of body weight administered as a drench or by dose syringe.

**FOR CHICKENS AND TURKEYS:** To arrive at the recommended dosages, use the enclosed contents of 2 scoops to prepare stock solutions as follows:

200 mg/gallon - dissolve 2 scoops (2.52 oz.) in 7,570 mL warm water (2 gallons)

400 mg/gallon - dissolve 2 scoops (2.52 oz.) in 3,785 mL warm water (1 gallon)

800 mg/gallon - dissolve 2 scoops (2.52 oz.) in 1,892 mL warm water (0.5 gallon)

These stock solutions should then be metered into drinking water at approximately 1 oz. per gallon.

At 200-800 mg/gallon two scoops will provide 63.8 to 255 gallons of medicated drinking water.

**SPECIAL DIRECTIONS FOR BABY CALVES AND BABY PIGS**

Administer this product one hour before or two hours after feeding milk or milk replacers. Provide clean (unmedicated) drinking water at all times.

**NOTE:** When using a watering trough, this product is to be administered twice a day in the drinking water of swine, calves and poultry. One-half of the recommended daily dosage level of antibiotic is to be consumed during each administration period, thus providing the drug in divided doses. The contents of this container (5 lb) will provide sufficient drug to treat 162,000 total pounds of swine or calves for a single day at the recommended dosage level of 10 mg/lb of body weight in divided doses. Or the contents of this container will treat 64,800 lbs of turkeys when supplied at 25 mg/lb or at 200-800 mg/gallon will provide 2,025 to 8,100 gallons of medicated drinking water for chickens or turkeys.

## RESTRICTED DRUG (California) - USE ONLY AS DIRECTED

## TAKE TIME OBSERVE LABEL DIRECTIONS

## CONTACT INFORMATION

To report suspected adverse drug events, for technical assistance, or to obtain a copy of the Safety Data Sheet (SDS), contact Dechra at (866) 933-2472. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at <http://www.fda.gov/reportanimalae>

TETRASOL is a trademark of Med-Pharmex, Inc.

## Manufactured for:

Dechra Veterinary Products, Overland Park, KS 66211 USA

**INDICATIONS:** For use in the control and treatment of the following conditions in swine, calves and poultry.

**SWINE AND CALVES**

**INDICATIONS:** Control and treatment of bacterial enteritis (scours) caused by *Escherichia coli*; bacterial pneumonia associated with *Actinobacillus pleuropneumoniae*, *Pasteurella* spp. and *Klebsiella* spp. sensitive to tetracycline hydrochloride.

**RECOMMENDED DOSAGE LEVEL:** Use soluble powder in the drinking water at a drug level of tetracycline hydrochloride per gallon to provide 10 mg/lb of body weight per day in divided doses.

**WARNING:**

**SWINE** - Do not slaughter animals for food purposes within 4 days of treatment.

**CALVES** - Do not slaughter animals for food purposes within 5 days of treatment.

A withdrawal period has not been established for this product in pre-ruminating calves.

Do not use in calves to be processed for veal.

**CAUTION:** Do not use for more than 5 consecutive days.

**DIRECTIONS FOR USE:** Administer for 3-5 days.

**CHICKENS**

**INDICATIONS:** Control of chronic respiratory disease (CRD) and air sac disease caused by *Mycoplasma gallisepticum* and *Escherichia coli*; infectious synovitis caused by *Mycoplasma synoviae* sensitive to tetracycline hydrochloride.

**RECOMMENDED DOSAGE LEVEL:** CRD and air sac disease: Use soluble powder in the drinking water at a drug level of 400-800 mg tetracycline hydrochloride per gallon. Infectious synovitis: Use soluble powder in the drinking water at a drug level of 200-400 mg tetracycline hydrochloride per gallon.

**TURKEYS**

**INDICATIONS:** Control of infectious synovitis caused by *Mycoplasma synoviae*; bluecomb (transmissible enteritis, coronavirus enteritis) complicated by organisms sensitive to tetracycline hydrochloride.

**RECOMMENDED DOSAGE LEVEL:** Infectious synovitis: Use soluble powder in the drinking water at a drug level of 400 mg tetracycline hydrochloride per gallon. Bluecomb: Use soluble powder in the drinking water at a drug level of tetracycline hydrochloride per gallon to provide 25 mg/lb of body weight per day in divided doses.

**WARNING:** Do not slaughter birds for food within 4 days of treatment.

Do not use for more than 14 consecutive days.

**DIRECTIONS FOR USE:** Administer for 7-14 days. Medicate at first clinical signs of disease or when experience indicates the disease may be a problem.

**RESTRICTED DRUG (California) - USE ONLY AS DIRECTED**

Lot:EXP  
Rev. February 2023

NDC 17033-434-05

**TETRASOL™**  
**SOLUBLE POWDER**  
(Tetracycline Hydrochloride Soluble Powder)  
**Antibiotic**

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**FOR USE IN ANIMALS ONLY  
KEEP OUT OF REACH OF CHILDREN**

Each pound contains 324 g of tetracycline hydrochloride.

Approved by FDA under ANADA # 200-234

**Net Weight:**  
**5 lbs. (2.27 kg)**

**Dechra**

**GENERAL CAUTION**

Prepare fresh solutions at least once a day. Solutions are not stable for more than 24 hours. Use as sole source of tetracycline. Diagnosis should be reconsidered if improvement is not noticed within 3 days. The concentration of drug required in medicated water must be adequate to compensate for variations in the age and class of animals, feed consumption, and environmental temperature and humidity, each of which affects water consumption.

**MIXING DIRECTIONS**

**FOR SWINE, CALVES AND TURKEYS:** 2.52 oz. (two scoops) dissolved in 500 mL (approximately 17 fl. oz.) of warm water will provide a stock solution of 100 mg of tetracycline hydrochloride activity per mL.

**FOR TURKEYS ONLY:** This stock solution when metered at approximately 1 oz. per gallon, will provide drinking water containing 2,957 mg of tetracycline hydrochloride activity per gallon.

The contents of two scoops are sufficient to deliver the recommended daily dosage levels as follows:

At 25 mg/lb of body weight ..... 2,040 total lbs of turkeys to be medicated.

At 10 mg/lb of body weight ..... 5,100 total lbs of swine or calves to be medicated.

Individual calf treatment ..... 5 mL (1 measuring teaspoonful) twice daily for each 100 lbs of body weight administered as a drench or by dose syringe.

**FOR CHICKENS AND TURKEYS:** To arrive at the recommended dosages, use the enclosed contents of 2 scoops to prepare stock solutions as follows:

200 mg/gallon - dissolve 2 scoops (2.52 oz.) in 7.570 mL warm water (2 gallons)

400 mg/gallon - dissolve 2 scoops (2.52 oz.) in 3.785 mL warm water (1 gallon)

800 mg/gallon - dissolve 2 scoops (2.52 oz.) in 1.892 mL warm water (0.5 gallon)

These stock solutions should then be metered into drinking water at approximately 1 oz. per gallon.

At 200-800 mg/gallon two scoops will provide 63.8 to 255 gallons of medicated drinking water.

**SPECIAL DIRECTIONS FOR BABY CALVES AND BABY PIGS**

Administer this product one hour before or two hours after feeding milk or milk replacers. Provide clean (unmedicated) drinking water at all times.

**NOTE:** When using a watering trough, this product is to be administered twice a day in the drinking water of swine, calves and poultry. One-half of the recommended daily dosage level of antibiotic is to be consumed during each administration period, thus providing the drug in divided doses. The contents of this container (5 lb) will provide sufficient drug to treat 162,000 total pounds of swine or calves for a single day at the recommended dosage level of 10 mg/lb of body weight in divided doses. Or the contents of this container will treat 64,800 lbs of turkeys when supplied at 25 mg/lb or at 200-800 mg/gallon will provide 2,025 to 8,100 gallons of medicated drinking water for chickens or turkeys.

**CONTACT INFORMATION**

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Dechra at (866) 933-2472.

For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at <http://www.fda.gov/reportanimalae>

TETRASOL is a trademark of Med-Pharmex, Inc.

**MANUFACTURED FOR:** Dechra Veterinary Products, Overland Park, KS 66211 USA

**TAKE TIME OBSERVE LABEL DIRECTIONS**

N 17033-434-05 5

## TETRASOL SOLUBLE POWDER

tetracycline hydrochloride powder, for solution

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:17033-434
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>TETRACYCLINE HYDROCHLORIDE</b> (UNII: P6R62377KV) (TETRACYCLINE - UNII:F8VB5M810T)	TETRACYCLINE HYDROCHLORIDE	714 g in 1 kg

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17033-434-05	2.27 kg in 1 CONTAINER		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200234	04/11/2023	

**Labeler** - Dechra Veterinary Products (362142734)

Revised: 4/2023

Dechra Veterinary Products